

# Noncommutativity, time and entropy bounds

Roberto Longo



*Pontifical Lateran University - November 27-28, 2018*

*IRAFS* workshop on **The origins and evolution of spacetime**

# Part I

## General introduction

## Noncommutativity and modular time

## Time as derived quantity

classical static space  $\rightarrow$  no time

quantum space  $\rightarrow$  quantum fluctuations

*no static quantum space may exist*

noncommutativity generates time

# The arrow of time

The arrow of time is viewed both classically and in quantum physics

thermodynamics → positive entropy

quantum mechanics → collapse of the wave function

Known question: is there a general frame to encompass both?

Of course, we keep in mind that time is a relative concept as we learnt from Einstein.

# Quantum Mechanics and Noncommutativity

*Schrödinger*

*Heisenberg*

*von Neumann uniqueness*



- Schrödinger:

$$i\hbar \frac{\partial}{\partial t} \psi(x, t) = H\psi(x, t)$$

Differential equations

- Heisenberg:

$$PQ - QP = i\hbar I$$

Linear operators on Hilbert space, **noncommutativity is essential!**

# Operator Algebras

$\mathcal{H}$  = Hilbert space,

$B(\mathcal{H})$  = algebra of all bounded linear operators on  $\mathcal{H}$ .

*Algebraic structure:* linear structure, multiplication:  $B(\mathcal{H})$  is a *\*algebra*

Derived structures:

*Order structure:*  $A \geq 0 \Leftrightarrow A = B^*B$ : *algebraic structure determines order structure*

*Metric structure:*

$\|A\|^2 = \inf\{\lambda > 0 : A^*A \leq \lambda I\}$ : *algebraic structure determines metric structure*

*$C^*$  property of the norm:*

$\|A^*A\| = \|A\|^2$ .  $B(\mathcal{H})$  is a  *$C^*$ -algebra*

## $C^*$ -algebras = noncommutative topology

**Gelfand-Naimark thm.**  $\exists$  contravariant functor  $F$  between category of *commutative*  $C^*$ -algebras and category of locally compact topological spaces:

$$\begin{array}{ccc} \mathfrak{A} & \xrightarrow{F} & \text{spec}(\mathfrak{A}) \\ \parallel & & \parallel \\ C(X) & \xleftarrow{F^{-1}} & X \end{array}$$

$C^*$ -algebra = dual of a topological space

Every  $C^*$ -algebra is isomorphic to a norm closed  $*$ -subalgebra of  $B(\mathcal{H})$ .

Noncommutative geometry =  $*$ -subalgebras of  $C^*$ -algebras  
+ structure (spectral triple), Connes NC geometry.

## von Neumann algebras = noncommutative measure theory

$\mathcal{M} \subset B(\mathcal{H})$  is a von Neumann algebra if  $\mathcal{M}$  is a  $*$ -algebra on  $\mathcal{H}$  and is weakly closed. Equivalently (von Neumann density theorem)

$$\mathcal{M} = \mathcal{M}''$$

with  $\mathcal{M}' = \{T \in B(\mathcal{H}) : TX = XT \quad \forall X \in \mathcal{M}\}$  the commutant.

$$\mathcal{M} \text{ abelian} \Leftrightarrow \mathcal{M} = L^\infty(X, \mu):$$

$$(\mathcal{M} = \{M_f : g \in L^2 \mapsto fg \in L^2\})$$

von Neumann algebra = dual of a measure space

Physics: *Observables* are selfadjoint elements  $X$  of  $\mathcal{M}$ , *states* are normalised positive linear functionals  $\varphi$ ,

$$\varphi(X) = \text{expected value of the observable } X \text{ in the state } \varphi$$



# Operator Algebras

Classical Commutative	Quantum Noncommutative
Manifold $X$ $C^\infty(X)$	*-algebra $A$
Topological space $X$ $C(X)$	$C^*$ -algebra $\mathfrak{A}$
Measure space $X$ $L^\infty(X, \mu)$	von Neumann algebra $\mathcal{A}$

# Quantum calculus with infinitely many degrees of freedom

CLASSICAL	Classical variables Differential forms Chern classes	Variational calculus Infinite dimensional manifolds Functions spaces Wiener measure
QUANTUM	Quantum geometry Fredholm operators Index Cyclic cohomology	Subfactors Bimodules, Endomorphisms Multiplicative index Supersymmetric QFT, $(\mathfrak{A}, \mathcal{H}, Q)$

# Thermal equilibrium states

A primary role in thermodynamics is played by the equilibrium distribution.

## Gibbs states

*Finite quantum system:*  $\mathfrak{A}$  matrix algebra with Hamiltonian  $H$  and evolution  $\tau_t = \text{Ad}e^{itH}$ . Equilibrium state  $\varphi$  at inverse temperature  $\beta$  is given by the Gibbs property

$$\varphi(X) = \frac{\text{Tr}(e^{-\beta H} X)}{\text{Tr}(e^{-\beta H})}$$

*What are the equilibrium states at infinite volume where there is no trace, no inner Hamiltonian?*


## KMS states (HHW, Baton Rouge conference 1967)

*Infinite volume.*  $\mathfrak{A}$  a  $C^*$ -algebra,  $\tau$  a one-par. automorphism group of  $\mathfrak{A}$ . A state  $\varphi$  of  $\mathfrak{A}$  is KMS at inverse temperature  $\beta > 0$  if for  $X, Y \in \mathfrak{A} \exists$  function  $F_{XY}$  s.t.

$$(a) F_{XY}(t) = \varphi(X\tau_t(Y))$$

$$(b) F_{XY}(t + i\beta) = \varphi(\tau_t(Y)X)$$

$F_{XY}$  bounded analytic on  $S_\beta = \{0 < \Im z < \beta\}$



*KMS states generalise Gibbs states, equilibrium condition for infinite systems*

## Tomita-Takesaki modular theory

$\mathcal{M}$  be a von Neumann algebra on  $\mathcal{H}$ ,  $\varphi = (\Omega, \cdot\Omega)$  normal faithful state on  $\mathcal{M}$ . Embed  $\mathcal{M}$  into  $\mathcal{H}$

$$\begin{array}{ccc} \mathcal{M} & \xrightarrow[\text{isometric}]{X \mapsto X^*} & \mathcal{M} \\ \downarrow X \mapsto X\Omega & & \downarrow X \mapsto X\Omega \\ \mathcal{H} & \xrightarrow[\text{non isometric}]{S_0: X\Omega \mapsto X^*\Omega} & \mathcal{H} \end{array}$$

$S = \bar{S}_0$ ,  $\Delta = S^*S > 0$  positive selfadjoint

$$t \in \mathbb{R} \mapsto \sigma_t^\varphi \in \text{Aut}(\mathcal{M})$$

$$\sigma_t^\varphi(X) = \Delta^{it} X \Delta^{-it}$$


*intrinsic dynamics associated with  $\varphi$  (modular automorphisms).*

# Modular theory and temperature

By a remarkable historical accident, Tomita announced the theorem at the 1967 Baton Rouge conference. Soon later Takesaki completed the theory and characterised the modular group by the KMS condition.

- $\sigma^\varphi$  is a **purely noncommutative** object (trivial in the commutative case)
- it is a **thermal equilibrium evolution** If  $\varphi(X) = \text{Tr}(\rho X)$  (type I case) then  $\sigma_t^\varphi(X) = \rho^{it} X \rho^{-it}$
- **arrow of modular time is thermodynamical** KMS condition at inverse temperature  $\beta = -1$
- **modular time is intrinsic modulo scaling** the rescaled group  $t \mapsto \sigma_{-t/\beta}^\varphi$  is physical,  $\beta^{-1}$  KMS temperature

# Time as thermodynamical effect




If time is the modular time, then the time arrow is associated both with positive entropy and with quantum structure!

## Jones index

Factors (von Neumann algebras with trivial center) are “very infinite-dimensional” objects. For an inclusion of factors  $\mathcal{N} \subset \mathcal{M}$  the Jones index  $[\mathcal{M} : \mathcal{N}]$  measure the relative size of  $\mathcal{N}$  in  $\mathcal{M}$ . Surprisingly, the index values are quantised:

$$[\mathcal{M} : \mathcal{N}] = 4 \cos^2\left(\frac{\pi}{n}\right), \quad n = 3, 4, \dots \quad \text{or} \quad [\mathcal{M} : \mathcal{N}] \geq 4$$

Jones index appears in many places in math and in physics.





# Quantum Field Theory

In QFT we have a quantum system with infinitely many degrees of freedom. The system is relativistic and there is particle creation and annihilation.

No mathematically rigorous QFT model with interaction still exists in 3+1 dimensions!

Haag local QFT:

$O$  spacetime regions  $\mapsto$  von Neumann algebras  $\mathcal{A}(O)$

to each region one associates the “noncommutative functions” with support in  $O$ .

## Local QFT nets


Local net  $\mathcal{A}$  on spacetime  $M$ : map  $O \subset M \mapsto \mathcal{A}(O) \subset B(\mathcal{H})$  s.t.

- *Isotony*,  $O_1 \subset O_2 \implies \mathcal{A}(O_1) \subset \mathcal{A}(O_2)$
- *Locality*,  $O_1, O_2$  spacelike  $\implies [\mathcal{A}(O_1), \mathcal{A}(O_2)] = \{0\}$
- *Poincaré covariance* (conformal, diffeomorphism) .
- *Positive energy and vacuum vector*.

$O \mapsto \mathcal{A}(O)$ : “Noncommutative chart” in QFT

# Bisognano-Wichmann theorem '75, Sewell comment '80

Rindler spacetime (wedge  $x_1 > 0$ ), vacuum modular group



$a$  uniform acceleration of  $\mathcal{O}$

$s/a$  proper time of  $\mathcal{O}$

$\beta = 2\pi/a$  inverse KMS temperature of  $\mathcal{O}$

Hawking-Unruh effect!

Time is geodesic, quantum gravitational effect!

# Representations

A (DHR) *representation*  $\rho$  of local net  $\mathcal{A}$  maps  $\mathcal{A}(O)$  on a different Hilbert space  $\mathcal{H}$  s.t. but  $\rho|_{\mathcal{A}(O')}$  is equivalent to the vacuum rep.

*Index-statistics theorem* (R.L. 1988):

$$d(\rho) = \left[ \mathcal{A}(O) : \rho(\mathcal{A}(O)) \right]^{\frac{1}{2}}$$

DHR dimension =  $\sqrt{\text{Jones index}}$

*Physical index*

*Anal. index*

(basis for a QFT index theorem).

Part II

Applications

Intrinsic bounds on entropy

## Bekenstein's bound

For decades, modular theory has played a central role in the operator algebraic approach to QFT, very recently several physical papers in other QFT settings are dealing with the modular group, although often in a heuristic (yet powerful) way!

I will discuss the Bekenstein bound, a universal limit on the entropy that can be contained in a physical system with given size and given total energy

If  $R$  is the radius of a sphere that can enclose our system, while  $E$  is its total energy including any rest masses, then its entropy  $S$  is bounded by

$$S \leq \lambda RE$$

The constant  $\lambda$  is often proposed  $\lambda = 2\pi$  (natural units).

## Casini's argument

Subtract to the bare entropy of the local state the entropy corresponding to the vacuum fluctuations.  $V$  bounded region.

The restriction  $\rho_V$  of a global state  $\rho$  to von Neumann algebra  $\mathcal{A}(V)$  has formally entropy given by

$$S(\rho_V) = -\text{Tr}(\rho_V \log \rho_V) ,$$

known to be infinite. So subtract the vacuum state entropy

$$S_V = S(\rho_V) - S(\rho_V^0)$$

with  $\rho_V^0$  the density matrix of the restriction of the vacuum state.

Similarly,  $K$  Hamiltonian for  $V$ , consider

$$K_V = \text{Tr}(\rho_V K) - \text{Tr}(\rho_V^0 K)$$

Bekenstein bound is now  $S_V \leq K_V$  which is equivalent to the **positivity of the relative entropy**

$$S(\rho_V | \rho_V^0) \equiv \text{Tr}(\rho_V (\log \rho_V - \log \rho_V^0)) \geq 0 ,$$

## Araki's relative entropy

An infinite quantum system is described by a von Neumann algebra  $\mathcal{M}$  typically not of type  $I$  so  $\text{Tr}$  does not exist; however Araki's relative entropy between two faithful normal states  $\varphi$  and  $\psi$  on  $\mathcal{M}$  is defined in general by

$$S(\varphi|\psi) \equiv -(\eta, \log \Delta_{\xi, \eta} \eta)$$

where  $\xi, \eta$  are cyclic vector representatives of  $\varphi, \psi$  and  $\Delta_{\xi, \eta}$  is the relative modular operator associated with  $\xi, \eta$ .

$$S(\varphi|\psi) \geq 0$$

positivity of the relative entropy

Relative entropy is one of the key concepts. We take the view that relative entropy is a primary concept and all entropy notions are derived concepts



## Analog of the Kac-Wakimoto formula (L. '97)

The root of our work relies in this formula for the incremental free energy of a black hole (cf. the Kac-Wakimoto formula, Kawahigashi, Xu, L.)

$H_\rho$  be the Hamiltonian for a uniformly accelerated observer in the Minkowski spacetime with acceleration  $a > 0$  in representation  $\rho$  (localised in the wedge for  $H_\rho$ )

$$(\Omega, e^{-tH_\rho}\Omega)|_{t=\beta} = d(\rho)$$

with  $\Omega$  the vacuum vector and  $\beta = \frac{2\pi}{a}$  the inverse Hawking-Unruh temperature.  $d(\rho)^2$  is Jones' index.

The left hand side is a generalised partition formula, so  $\log d(\rho)$  has an **entropy meaning** in accordance with Pimsner-Popa work.

Here we generalise this formula

## CP maps, quantum channels and entropy

$\mathcal{N}, \mathcal{M}$  vN algebras. A linear map  $\alpha : \mathcal{N} \rightarrow \mathcal{M}$  is completely positive if

$$\alpha \otimes \text{id}_n : \mathcal{N} \otimes \text{Mat}_n(\mathbb{C}) \rightarrow \mathcal{M} \otimes \text{Mat}_n(\mathbb{C})$$

is positive  $\forall n$  (quantum operation)

$\omega$  faithful normal state of  $\mathcal{M}$  and  $\alpha : \mathcal{N} \rightarrow \mathcal{M}$  CP map as above.  
Set

$$H_\omega(\alpha) \equiv \sup_{(\omega_i)} \sum_j S(\omega|\omega_j) - S(\omega \cdot \alpha|\omega_j \cdot \alpha)$$

supremum over all  $\omega_j$  with  $\sum_j \omega_j = \omega$ .

The **conditional entropy**  $H(\alpha)$  of  $\alpha$  is defined by

$$H(\alpha) = \inf_{\omega} H_\omega(\alpha)$$

infimum over all “full” states  $\omega$  for  $\alpha$ . Clearly  $H(\alpha) \geq 0$  because  $H_\omega(\alpha) \geq 0$  by the **monotonicity of the relative entropy**.  
 $\alpha$  is a **quantum channel** if its conditional entropy  $H(\alpha)$  is finite.

## Generalisation of Stinespring dilation

Let  $\alpha : \mathcal{N} \rightarrow \mathcal{M}$  be a normal, completely positive unital map between the vN algebras  $\mathcal{N}$ ,  $\mathcal{M}$ . A pair  $(\rho, v)$   $\rho : \mathcal{N} \rightarrow \mathcal{M}$  a homomorphism,  $v \in \mathcal{M}$  an isometry s.t.

$$\alpha(n) = v^* \rho(n) v, \quad n \in \mathcal{N}.$$

$(\rho, v)$  is *minimal* if the left support of  $\rho(\mathcal{N})v\mathcal{H}$  is equal to 1.

**Thm** Let  $\alpha : \mathcal{N} \rightarrow \mathcal{M}$  be a normal, CP unital map with  $\mathcal{N}$ ,  $\mathcal{M}$  properly infinite. There exists a minimal dilation pair  $(\rho, v)$  for  $\alpha$ . If  $(\rho_1, v_1)$  is another minimal pair,  $\exists!$  unitary  $u \in \mathcal{M}$  such that

$$u\rho(n) = \rho_1(n)u, \quad v_1 = uv, \quad n \in \mathcal{N}$$

We have

$$H(\alpha) = \log \text{Ind}(\alpha) \quad (\text{minimal index})$$

# Bimodules and CP maps

Let  $\alpha : \mathcal{N} \rightarrow \mathcal{M}$  be a completely positive, normal, unital map and  $\omega$  a faithful normal state of  $\mathcal{M}$

$\exists!$   $\mathcal{N} - \mathcal{M}$  bimodule  $\mathcal{H}_\alpha$ , with a cyclic vector  $\xi_\alpha \in \mathcal{H}$  and left and right actions  $\ell_\alpha$  and  $r_\alpha$ , such that

$$(\xi_\alpha, \ell_\alpha(n)\xi_\alpha) = \omega_{\text{out}}(n), \quad (\xi_\alpha, r_\alpha(m)\xi_\alpha) = \omega_{\text{in}}(m),$$

with  $\omega_{\text{in}} \equiv \omega$ ,  $\omega_{\text{out}} \equiv \omega_{\text{in}} \cdot \alpha$ . Converse is true.

CP map  $\alpha \longleftrightarrow$  cyclic bimodule  $\mathcal{H}_\alpha$

We have

$$H(\alpha) = \log \text{Ind}(\mathcal{H}_\alpha) \quad (\text{Jones' index})$$

# Promoting modular theory to the bimodule setting

$\mathcal{H}$  an  $\mathcal{N} - \mathcal{M}$ -bimodule with finite Jones' index  $\text{Ind}(\mathcal{H})$

Given faithful, normal, states  $\varphi, \psi$  on  $\mathcal{N}$  and  $\mathcal{M}$ , I define the **modular operator**  $\Delta_{\mathcal{H}}(\varphi|\psi)$  of  $\mathcal{H}$  with respect to  $\varphi, \psi$  as

$$\Delta_{\mathcal{H}}(\varphi|\psi) \equiv d(\varphi \cdot \ell^{-1})/d(\psi \cdot r^{-1} \cdot \varepsilon) ,$$

Connes' spatial derivative,  $\varepsilon : \ell(\mathcal{N})' \rightarrow r(\mathcal{M})$  is the minimal conditional expectation

$\log \Delta_{\mathcal{H}}(\varphi|\psi)$  is called the **modular Hamiltonian** of the bimodule  $\mathcal{H}$ , or of the quantum channel  $\alpha$  if  $\mathcal{H}$  is associated with  $\alpha$ .

# Properties of the modular Hamiltonian

If  $\mathcal{N}$ ,  $\mathcal{M}$  factors

$$\Delta_{\mathcal{H}}^{it}(\varphi|\psi)\ell(n)\Delta_{\mathcal{H}}^{-it}(\varphi|\psi) = \ell(\sigma_t^\varphi(n))$$

$$\Delta_{\mathcal{H}}^{it}(\varphi|\psi)r(m)\Delta_{\mathcal{H}}^{-it}(\varphi|\psi) = r(\sigma_t^\psi(m))$$

(implements the dynamics)

$$\Delta_{\mathcal{H}}^{it}(\varphi_1|\varphi_2) \otimes \Delta_{\mathcal{K}}^{it}(\varphi_2|\varphi_3) = \Delta_{\mathcal{H} \otimes \mathcal{K}}^{it}(\varphi_1|\varphi_3)$$

(additivity of the energy)

$$\Delta_{\mathcal{H}}^{it}(\varphi_2|\varphi_1) = d_{\mathcal{H}}^{-i2t} \overline{\Delta_{\mathcal{H}}^{it}(\varphi_1|\varphi_2)}$$

If  $T : \mathcal{H} \rightarrow \mathcal{H}'$  is a bimodule intertwiner, then

$$T \Delta_{\mathcal{H}}^{it}(\varphi_1|\varphi_2) = (d_{\mathcal{H}'}/d_{\mathcal{H}})^{it} \Delta_{\mathcal{H}'}^{it}(\varphi_1|\varphi_2) T$$

Connes's bimodule tensor product w.r.t.  $\varphi_2$ ;  $d_{\mathcal{H}} = \sqrt{\text{Ind}(\mathcal{H})}$

# Physical Hamiltonian

We may modify the modular Hamiltonian in order to fulfil the right physical requirements (additivity of energy, invariance under charge conjugation,...)

$$K(\varphi_1|\varphi_2) = -\log \Delta_{\mathcal{H}}(\varphi_1|\varphi_2) - \log d$$

is the **physical Hamiltonian** (at inverse temperature 1).

The physical Hamiltonian at inverse temperature  $\beta > 0$  is given by

$$-\beta^{-1} \log \Delta - \beta^{-1} \log d$$

From the modular Hamiltonian to the physical Hamiltonian:

$$-\log \Delta \xrightarrow{\text{shifting}} -\log \Delta - \log d \xrightarrow{\text{scaling}} \beta^{-1} (-\log \Delta - \log d)$$

The shifting is **intrinsic**, the scaling is to be determined by the context!

# Modular and Physical Hamiltonians for a quantum channel

We now are going to compare two states of a physical system,  $\omega_{\text{in}}$  is a suitable reference state, e.g. the vacuum in QFT, and  $\omega_{\text{out}}$  is a state that can be reached from  $\omega_{\text{in}}$  by some physically realisable process (quantum channel).

$\alpha : \mathcal{N} \rightarrow \mathcal{M}$  be a quantum channel (normal, unital CP map with finite entropy) and  $\omega_{\text{in}}$  a faithful normal state of  $\mathcal{M}$ .  $\omega_{\text{out}} = \omega_{\text{in}} \cdot \alpha$

$$\log \Delta_{\alpha} \equiv \log \Delta_{\mathcal{H}_{\alpha}}$$

$$K_{\alpha} = \beta^{-1} K_{\mathcal{H}_{\alpha}} = \beta^{-1} ( - \log \Delta_{\mathcal{H}_{\alpha}} - \log d_{\mathcal{H}_{\alpha}} )$$

(physical Hamiltonian at inverse temperature  $\beta$ )

$K_{\alpha}$  may be considered as a local Hamiltonian associated with  $\alpha$  and the state transfer with input state  $\omega_{\text{in}}$ .



# Thermodynamical quantities

The **entropy**  $S \equiv S_{\alpha, \omega_{\text{in}}}$  of  $\alpha$  is

$$S = -(\hat{\xi}, \log \Delta_{\alpha} \hat{\xi})$$

where  $\hat{\xi}$  is a vector representative of the state  $\omega_{\text{in}} \cdot r^{-1} \cdot \varepsilon$  in  $\mathcal{H}_{\alpha}$ .

The quantity

$$E = (\hat{\xi}, K \hat{\xi})$$

is the **relative energy** w.r.t. the states  $\omega_{\text{in}}$  and  $\omega_{\text{out}}$ .

The **free energy**  $F$  is now defined by the relative partition function

$$F = -\beta^{-1} \log(\hat{\xi}, e^{-\beta K} \hat{\xi})$$

$F$  satisfies the **thermodynamical relation**

$$F = E - TS$$

## A form of Bekenstein bound

As  $F = \frac{1}{2}\beta^{-1}H(\alpha)$ , we have

$$F \geq 0 \quad (\text{positivity of the free energy})$$

because

$$H(\alpha) \geq 0 \quad (\text{monotonicity of the entropy})$$

So the above thermodynamical relation

$$F = E - \beta^{-1}S$$

entails the following general, rigorous version of the Bekenstein bound

$$S \leq \beta E$$

To determine  $\beta$  we have to plug this general formula in a physical context

## Fixing the temperature in QFT

$O$  a spacetime region s.t. the modular group  $\sigma_t^\omega$  of the local von Neumann algebra  $\mathcal{A}(O)$  associated with vacuum  $\omega$  has a geometric meaning. So there is a geometric flow  $\theta_s : O \rightarrow O$  and a re-parametrisation of  $\sigma_t^\omega$  that acts covariantly w.r.t  $\theta$ .

Motivated by the Rindler case one define locally the inverse temperature by

$$\beta_s = \left\| \frac{d\theta_s}{ds} \right\|$$

the Minkowskian length of the tangent vector to the modular orbit. Namely  $d\tau = \beta_s ds$  with  $\tau$  proper time (cf. Connes and Rovelli).

## Schwarzschild black hole

Schwarzschild-Kruskal spacetime of mass  $M > 0$ , namely the region inside the event horizon, and  $\mathcal{N} \equiv \mathcal{A}(O)$  the local von Neumann algebra associated with  $O$  on the underlying Hilbert space  $\mathcal{H}$ ,  $O$  Schwarzschild black hole region,  $\omega$  vacuum state

$\mathcal{H}$  is a  $\mathcal{N} - \mathcal{N}$  bimodule, indeed the identity  $\mathcal{N} - \mathcal{N}$  bimodule  $L^2(\mathcal{N})$  associated with  $\omega$ .

The modular group of  $\mathcal{A}(O)$  associated with  $\omega$  is geometric and corresponds to the geodesic flow. KMS Hawking temperature is

$$T = 1/8\pi M = 1/4\pi R$$

with  $R = 2M$  the Schwarzschild radius, then

$$S \leq 4\pi RE$$

with  $S$  the entropy associated with the state transfer of  $\omega$  by a quantum channel, and  $E$  the corresponding relative energy.

## Conformal QFT

Conformal Quantum Field Theory on the Minkowski spacetime, any spacetime dimension.  $O_R$  double cone with basis a radius  $R > 0$  sphere centered at the origin and  $\mathcal{A}(O_R)$  associated local vN algebra.

The modular group of  $\mathcal{A}(O_R)$  w.r.t. the vacuum state  $\omega$  has a geometrical meaning (Hislop, L. 1982):

$$\Delta_{O_R}^{-is} = U(\Lambda_{O_R}(2\pi s))$$

with  $U$  is the representation of the conformal group and  $\Lambda_{O_R}$  is a one-parameter group of conformal transformation leaving  $O_R$  globally invariant and conjugate to the boost one-parameter group of pure Lorentz transformations.

The inverse temperature  $\beta_R = \left\| \frac{d}{ds} \Lambda_{O_R}(s) \mathbf{x} \right\|_{s=0}$  in  $O_R$  is maximal on the time-zero basis of  $O_R$ , in fact at the origin  $\mathbf{x} = \mathbf{0}$  with value

$$\beta_R = \pi R$$

So

$$S \leq \pi R E$$

with  $S$  and  $E$  the entropy and energy associated with any quantum channel by the vacuum state.

## Summary

von Neumann algebra  $\longleftrightarrow$  quantum system

CP map with finite entropy between q. systems  $\longleftrightarrow$  quantum channel

quantum channel  $\longleftrightarrow$  finite index bimodule

finite index bimodule and state  $\longrightarrow$  modular Hamiltonian

modular Hamiltonian & physical functoriality  $\longrightarrow$  phys. Hamiltonian

modular and physical Hamiltonians  $\longrightarrow F = E - TS$

$F = E - TS$  & autom. positivity of the free energy  $F \longrightarrow S \leq \beta E$

$S \leq \beta E$  & geometrical modular flow  $\longrightarrow$  Bekenstein's bound

## Landauer's bound for infinite systems

Let  $\alpha : \mathcal{N} \rightarrow \mathcal{M}$  be a quantum channel between quantum systems  $\mathcal{N}$ ,  $\mathcal{M}$ . If  $\alpha$  is irreversible, then

$$F_\alpha \geq \frac{1}{2} kT \log 2$$

The original lower bound for the incremental free energy is  $F_\alpha \geq kT \log 2$ , it remains true for finite-dimensional systems  $\mathcal{N}$ ,  $\mathcal{M}$ .



## Entropy distribution of localised states

Case of  $U(1)$ -current  $j$ :  $\ell$  real function in  $S(\mathbb{R})$  and  $t \in \mathbb{R}$ . We have

$$S(t) = \pi \int_t^{+\infty} (x - t) \ell^2(x) dx ,$$

$S(t)$  vacuum relative entropy of excited state by  $j \mapsto j + \ell$ , so

$$S'(t) = -\pi \int_t^{+\infty} \ell^2(x) dx \leq 0 ,$$

$$S''(t) = \pi \ell^2(t) \geq 0$$


positivity of  $S''$

## Quantum Null Energy Condition

The vacuum energy density is  $E(t) = \frac{1}{2}\ell^2(t)$  so we have here the QNEC:

$$E(t) = \frac{1}{2\pi} S''(t) \geq 0$$

QNEC is not saturated in every point of positive energy density.



**Figure:** Two distributions, blue and red, for the same charge  $q = \int \ell$ . The dashed lines plot the corresponding entropy density rate  $S''(t)$ : blue high entropy, red low entropy.